



Measles

Measles *Second dose should be given at least four weeks after the first dose. Measles, Mumps and Rubella (MMR) combined vaccine is preferred for those who may also be susceptible to mumps and/or rubella.*

Dose 1 _____

Dose 2 _____

Mumps

Mumps *One dose only, no booster. MMR combined vaccine is preferred for those who may also be susceptible to measles and/or rubella.*

Vaccine _____

Rubella

Rubella *One dose only, no booster. MMR combined vaccine is preferred for those who may also be susceptible to measles and/or mumps.*

Vaccine _____

Varicella zoster (Chickenpox)

Varicella zoster *Adults should receive second dose four to eight weeks after first dose. May be cost-effective to test for immunity before vaccinating.*

Dose 1 _____

Dose 2 _____



Influenza

Influenza	Single dose given annually in the fall or early winter months. Healthcare workers who should be vaccinated include those who are or care for patients who are at high-risk for complications from influenza. People in the high-risk group are those 65 years or older; pregnant women who will be in their second or third trimester during the influenza season; people with chronic pulmonary or cardiovascular diseases; people who have required medical follow-up or hospitalization within the past year because of chronic metabolic disease (such as diabetes), renal dysfunction, hemoglobinopathies, or immunosuppression.
------------------	---

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____

Vaccine _____